TB Disinfectant

Ready to Use



Disinfection is a process, not a product.

Step 1: Clean, Step 2: <u>Disinfect</u>*



- Our unique 2 product system provides powerful cleaning and broad-spectrum disinfection to reduce the risk of hazardous crosscontamination on treated surfaces!
- A perfect disinfectant to pair with bloodborne pathogen clean-up kits.
- Meets OSHA Bloodborne Pathogen Standard for HIV, HBV & HCV.

- Kills Norovirus, Poliovirus Type 1, Tuberculosis, MRSA and VRE.**
- Contains no abrasives or bleach.
- Contains no phosphates or phosphorous.
- Destroys mold and mildew odors at the source.

EnvirOx's TB Disinfectant is designed as part of a 2 product system. First, clean the surface with EnvirOx Heavy Duty RED® dilution from one of our proprietary hydrogen peroxide based multi-purpose cleaners. Then, disinfect the area using TB Disinfectant RTU as directed on the product label.

Use to disinfect all major touch points around your facility (i.e., door handles, faucets, telephones, desk/tabletops) to provide a healthier environment for your staff, students or customers.

Use as:

- Disinfectant
- Tuberculocidal
- Virucide
- Fungicide
- Midewstat
- Deodorizer

STEP 1: CLEAN



STEP 2: DISINFECT



Kills SARS-CoV-2 (COVID-19 virus) on hard non-porous surfaces in just 60 seconds!

TB Disinfectant RTU

Description	Disinfectant, Fungicide, Mildewstat, Virucide, Tuberculocidal, Deodorizer	
Fragrance	Lemon	
Packaging	12 x 32 oz Bottles	
Dispensing	Ready-to-Use (RTU)	
Yield Per Case	12 Quarts	
Certification	EPA: 1839-83-69268	
Shelf Life	1 Year	

BACTERICIDAL ACTIVITY: Using approved AOAC test methods in the presence of 5% blood serum and a **3** minute contact time, unless otherwise noted, this product kills the following organisms on hard non-porous inanimate surfaces:

Organism	Contact Time
Pseudomonas aeruginosa	3 minutes
Staphylococcus aureus	3 minutes
Salmonella (choleraesuis) enterica	3 minutes
Escherichia coli	3 minutes
Escherichia coli 0157:H7	3 minutes
Streptococcus pyogenes (Necrotizing Fasciitis-Group A) (V.A. Medical Center Isolate 04001)	3 minutes
Listeria monocytogenes	3 minutes
Yersinia enterocolitica	3 minutes
Enterococcus faecium	3 minutes
Corynebacterium ammoniagenes	3 minutes
Color on the Armely contribute	3 minutes
Salmonella (typhi) enterica	
ANTIBIOTIC-RESISTANT: Using approved AOAC test methods in the presence minute contact time, unless otherwise noted, this product kills the following or inanimate surfaces:	of 5% blood serum and a 3
ANTIBIOTIC-RESISTANT: Using approved AOAC test methods in the presence minute contact time, unless otherwise noted, this product kills the following or	of 5% blood serum and a 3
ANTIBIOTIC-RESISTANT: Using approved AOAC test methods in the presence minute contact time, unless otherwise noted, this product kills the following o inanimate surfaces:	of 5% blood serum and a 3 rganisms on hard non-porous
ANTIBIOTIC-RESISTANT: Using approved AOAC test methods in the presence minute contact time, unless otherwise noted, this product kills the following or inanimate surfaces: Methicillin resistant Staphylococcus aureus (MRSA)	of 5% blood serum and a 3 rganisms on hard non-porous
ANTIBIOTIC-RESISTANT: Using approved AOAC test methods in the presence minute contact time, unless otherwise noted, this product kills the following o inanimate surfaces: Methicillin resistant Staphylococcus aureus (MRSA) Vancomycin resistant Enterococcus faecalis (VRE)	of 5% blood serum and a 3 rganisms on hard non-porous 3 minutes 3 minutes

EMERGING VIRAL PATHOGEN CLAIMS

This product qualifies for emerging viral pathogen claims per the EPA's 'Guidance to Registrants: Process for Making Claims Against Emerging Viral Pathogens not on EPA-Registered Disinfectant Labels' when used in accordance with the appropriate use directions indicated below:

This product meets the criteria to make claims against certain emerging viral pathogens from the following viral categories:

- · Enveloped viruses
- · Large non-enveloped viruses
- Small non-enveloped viruses


For an emerging viral pathogen that is a/an	follow the directions for use for the following organisms on the label:
Enveloped virus	Human Rotavirus Rhinovirus Type 39 Hepatitis A Virus Poliovirus Type 1 Canine Parvovirus
Large non-enveloped virus	Rhinovirus Type 39 Hepatitis A Virus Poliovirus Type 1 Canine Parvovirus
Small non-enveloped virus §	Hepatitis A Virus § Poliovirus Type 1 § Canine Parvovirus §

§ The directions for use for the supporting viruses with the longest contact time should be used for small non-enveloped emerging viruses. In this case, 10 minutes for the indicated viruses.

VIRUCIDIAL ACTIVITY:	Contact Time
Virus Enveloped	
HIV-1 (associated with AIDS)	1 minute
Avian Influenza A strain H3N2	2 minutes
Avian Influenza A strain H9N2	2 minutes
Hepatitus B Virus (HBV)	1 minute
Hepatitis C Virus (HCV)	1 minute
Duck Hepatitis B Virus (DHBV)	1 minute
Bovine Viral Diarrhea Virus (BVDV)	1 minute
Human Coronavirus	2 minutes
SARS associated Coronavirus	2 minutes
SARS-Related Coronavirus 2 [SARS-CoV-2] USA-WA1/2020 BEI NR-52281	1 minute
Rabies Virus	30 seconds
Paramyxovirus (Mumps)	3 minutes
Virus Large Non-Enveloped	
Rotavirus	3 minutes
Virus Small Non-Enveloped	
Norovirus (Norwalk virus)	30 seconds
Feline Calicivirus	30 seconds
Canine Parvovirus	10 minutes
Poliovirus Type 1	10 minutes
Rhinovirus type 39	3 minutes
Hepatitis A Virus (HAV) (University of Ottawa)	10 minutes
MILDEW / FUNGI:	Contact Time
Trichophyton mentagrophytes (athlete's foot fungus)	10 minutes



Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA) (NRS 384) Genotype USA300



3 minutes

1-800-281-9604 • www.enviroxclean.com enviroxinfo@enviroxclean.com

